

Continuous-Time Fourier Transform



Content

- Introduction
- Fourier Integral
- Fourier Transform
- Properties of Fourier Transform
- Convolution
- Parseval's Theorem

Continuous-Time Fourier Transform


Introduction

The Topic

	Continuous Time	Discrete Time
Periodic	Fourier Series	Discrete Fourier Transform
Aperiodic	Continuous Fourier Transform	Fourier Transform

Review of Fourier Series

- Deal with continuous-time periodic signals.
- Discrete frequency spectra.



Two Forms for Fourier Series

Sinusoidal
Form

$$f(t) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{2\pi n t}{T} + \sum_{n=1}^{\infty} b_n \sin \frac{2\pi n t}{T}$$

$$a_0 = \frac{2}{T} \int_{-T/2}^{T/2} f(t) dt$$

$$a_n = \frac{2}{T} \int_{-T/2}^{T/2} f(t) \cos n\omega_0 t dt$$

$$b_n = \frac{2}{T} \int_{-T/2}^{T/2} f(t) \sin n\omega_0 t dt$$


Complex
Form:

$$f(t) = \sum_{n=-\infty}^{\infty} c_n e^{jn\omega_0 t}$$

$$c_n = \frac{1}{T} \int_{-T/2}^{T/2} f(t) e^{-jn\omega_0 t} dt$$

How to Deal with Aperiodic Signal?

A Periodic Signal



If $T \rightarrow \infty$, what happens?

Continuous-Time Fourier Transform

Fourier Integral

Fourier Integral

$$f_T(t) = \sum_{n=-\infty}^{\infty} c_n e^{jn\omega_0 t} \quad c_n = \frac{1}{T} \int_{-T/2}^{T/2} f_T(t) e^{-jn\omega_0 t} dt$$

$$= \sum_{n=-\infty}^{\infty} \left[\frac{1}{T} \int_{-T/2}^{T/2} f_T(\tau) e^{-jn\omega_0 \tau} d\tau \right] e^{jn\omega_0 t}$$

$$\omega_0 = \frac{2\pi}{T} \quad \rightarrow \quad \frac{1}{T} = \frac{\omega_0}{2\pi}$$

$$= \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} \left[\int_{-T/2}^{T/2} f_T(\tau) e^{-jn\omega_0 \tau} d\tau \right] \omega_0 e^{jn\omega_0 t}$$

$$\text{Let } \Delta\omega = \omega_0 = \frac{2\pi}{T}$$

$$= \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} \left[\int_{-T/2}^{T/2} f_T(\tau) e^{-jn\omega_0 \tau} d\tau \right] e^{jn\omega_0 t} \Delta\omega$$

$$T \rightarrow \infty \Rightarrow d\omega = \Delta\omega \approx 0$$

$$= \frac{1}{2\pi} \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} f_T(\tau) e^{-j\omega \tau} d\tau \right] e^{j\omega t} d\omega$$

Fourier Integral

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \underbrace{\left[\int_{-\infty}^{\infty} f(\tau) e^{-j\omega\tau} d\tau \right]}_{F(j\omega)} e^{j\omega t} d\omega$$

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(j\omega) e^{j\omega t} d\omega \quad \text{Synthesis}$$

$$F(j\omega) = \int_{-\infty}^{\infty} f(t) e^{-j\omega t} dt \quad \text{Analysis}$$

Fourier Series vs. Fourier Integral

Fourier Series:

$$f(t) = \sum_{n=-\infty}^{\infty} c_n e^{jn\omega_0 t}$$

Period Function

$$c_n = \frac{1}{T} \int_{-T/2}^{T/2} f_T(t) e^{-jn\omega_0 t} dt$$

Discrete Spectra

Fourier Integral:

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(j\omega) e^{j\omega t} d\omega$$

Non-Period Function

$$F(j\omega) = \int_{-\infty}^{\infty} f(t) e^{-j\omega t} dt$$

Continuous Spectra

Continuous-Time Fourier Transform

Fourier Transform



Fourier Transform Pair

Inverse Fourier Transform:

$$f(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(j\omega) e^{j\omega t} d\omega$$

Synthesis

Fourier Transform:

$$F(j\omega) = \int_{-\infty}^{\infty} f(t) e^{-j\omega t} dt$$

Analysis

Existence of the Fourier Transform

Sufficient Condition:

$f(t)$ is absolutely integrable, i.e.,


$$\int_{-\infty}^{\infty} |f(t)| dt < \infty$$

Continuous Spectra


$$F(j\omega) = \int_{-\infty}^{\infty} f(t)e^{-j\omega t} dt$$

$$F(j\omega) = F_R(j\omega) + jF_I(j\omega)$$

$$= \underbrace{|F(j\omega)|}_{\text{Magnitude}} e^{j\underbrace{\phi(\omega)}_{\text{Phase}}}$$




Example




$$\begin{aligned} F(j\omega) &= \int_{-\infty}^{\infty} f(t)e^{-j\omega t} dt = \int_{-1}^1 e^{-j\omega t} dt = \frac{1}{-j\omega} e^{-j\omega t} \Big|_{-1}^1 \\ &= \frac{j}{\omega} (e^{-j\omega} - e^{j\omega}) = \frac{2\sin \omega}{\omega} = 2\text{sinc } 2\pi f \end{aligned}$$

Example




$$F(j\omega) = \int_{-\infty}^{\infty} f(t)e^{-j\omega t} dt$$

$$= \frac{j}{\omega} (e^{-j\omega} - e^{j\omega})$$




Example



$$\begin{aligned} F(j\omega) &= \int_{-\infty}^{\infty} f(t) e^{-j\omega t} dt = \int_0^{\infty} e^{-\alpha t} e^{-j\omega t} dt \\ &= \int_0^{\infty} e^{-(\alpha + j\omega)t} dt = \frac{1}{\alpha + j\omega} \end{aligned}$$

Example



$$F(j\omega) = \int_{-\infty}^{\infty} f$$
$$= \int_0^{\infty} e^{-$$

Continuous-Time Fourier Transform

Properties of
Fourier Transform

Notation

$$\mathcal{F}[f(t)] = F(j\omega)$$

$$\mathcal{F}^{-1}[F(j\omega)] = f(t)$$



$$f(t) \xleftrightarrow{\mathcal{F}} F(j\omega)$$

Linearity

$$a_1 f_1(t) + a_2 f_2(t) \xleftrightarrow{\mathcal{F}} a_1 F_1(j\omega) + a_2 F_2(j\omega)$$

Proved by yourselves

Time Scaling

$$f(at) \xleftrightarrow{F} \frac{1}{|a|} F\left(j\frac{\omega}{a}\right)$$

Proved by yourselves

Time Reversal

$$f(-t) \xleftrightarrow{F} F(-j\omega)$$

Pf)

$$\begin{aligned} F[f(-t)] &= \int_{-\infty}^{\infty} f(-t)e^{-j\omega t} dt = \int_{t=-\infty}^{t=\infty} f(-t)e^{-j\omega t} dt \\ &= \int_{-t=-\infty}^{-t=\infty} f(t)e^{j\omega t} d(-t) = \int_{-t=-\infty}^{-t=\infty} f(t)e^{j\omega t} d(-t) \\ &= -\int_{t=\infty}^{t=-\infty} f(t)e^{j\omega t} dt = \int_{t=-\infty}^{t=\infty} f(t)e^{j\omega t} dt \\ &= \int_{-\infty}^{\infty} f(t)e^{j\omega t} dt = F(-j\omega) \end{aligned}$$

Time Shifting

$$f(t - t_0) \xleftrightarrow{F} F(j\omega)e^{-j\omega t_0}$$

$$\begin{aligned} \text{Pf)} \quad F[f(t - t_0)] &= \int_{-\infty}^{\infty} f(t - t_0)e^{-j\omega t} dt = \int_{t=-\infty}^{t=\infty} f(t - t_0)e^{-j\omega t} dt \\ &= \int_{t+t_0=-\infty}^{t+t_0=\infty} f(t)e^{-j\omega(t+t_0)} d(t + t_0) \\ &= e^{-j\omega t_0} \int_{t=-\infty}^{t=\infty} f(t)e^{-j\omega t} dt \\ &= e^{-j\omega t_0} \int_{-\infty}^{\infty} f(t)e^{-j\omega t} dt = F(j\omega)e^{-j\omega t_0} \end{aligned}$$

Frequency Shifting (Modulation)

$$f(t)e^{j\omega_0 t} \xleftrightarrow{F} F[j(\omega - \omega_0)]$$

Pf)

$$\begin{aligned} F[f(t)e^{j\omega_0 t}] &= \int_{-\infty}^{\infty} f(t)e^{j\omega_0 t} e^{-j\omega t} dt \\ &= \int_{-\infty}^{\infty} f(t)e^{-j(\omega - \omega_0)t} dt \\ &= F[j(\omega - \omega_0)] \end{aligned}$$

Symmetry Property

$$\mathcal{F}[F(jt)] = 2\pi f(-\omega)$$

Pf)

$$2\pi f(t) = \int_{-\infty}^{\infty} F(j\omega)e^{j\omega t} d\omega$$

$$2\pi f(-t) = \int_{-\infty}^{\infty} F(j\omega)e^{-j\omega t} d\omega$$

Interchange symbols ω and t

$$2\pi f(-\omega) = \int_{-\infty}^{\infty} F(jt)e^{-j\omega t} dt = \mathcal{F}[F(jt)]$$

Fourier Transform for Real Functions

If $f(t)$ is a real function, and $F(j\omega) = F_R(j\omega) + jF_I(j\omega)$

 $F(-j\omega) = F^*(j\omega)$

$$F(j\omega) = \int_{-\infty}^{\infty} f(t)e^{-j\omega t} dt$$

$$F^*(j\omega) = \int_{-\infty}^{\infty} f(t)e^{j\omega t} dt = F(-j\omega)$$

Fourier Transform for Real Functions

If $f(t)$ is a real function, and $F(j\omega) = F_R(j\omega) + jF_I(j\omega)$

➔ $F(-j\omega) = F^*(j\omega)$

➔ $F_R(j\omega)$ is even, and $F_I(j\omega)$ is odd.

$$\underbrace{F_R(-j\omega)} = \underbrace{F_R(j\omega)} \quad \underbrace{F_I(-j\omega)} = \underbrace{-F_I(j\omega)}$$

➔ *Magnitude spectrum* $|F(j\omega)|$ is even, and *phase spectrum* $\phi(\omega)$ is odd.

Fourier Transform for Real Functions

If $f(t)$ is real and even

→ $F(j\omega)$ is real ✓

Pf)

Even → $f(t) = f(-t)$

→ $F(j\omega) = F(-j\omega)$

Real → $F(-j\omega) = F^*(j\omega)$

→ $F(j\omega) = F^*(j\omega)$

If $f(t)$ is real and odd

→ $F(j\omega)$ is pure imaginary ✓

Pf)

Odd → $f(t) = -f(-t)$

→ $F(j\omega) = -F(-j\omega)$

Real → $F(-j\omega) = F^*(j\omega)$

→ $F(j\omega) = -F^*(j\omega)$

Example:

$$\mathcal{F}[f(t)] = F(j\omega)$$


$$\mathcal{F}[f(t) \cos \omega_0 t] = ?$$

Sol)

$$f(t) \cos \omega_0 t = \frac{1}{2} f(t) (e^{j\omega_0 t} + e^{-j\omega_0 t})$$

$$\begin{aligned} \mathcal{F}[f(t) \cos \omega_0 t] &= \frac{1}{2} \mathcal{F}[f(t) e^{j\omega_0 t}] + \frac{1}{2} \mathcal{F}[f(t) e^{-j\omega_0 t}] \\ &= \frac{1}{2} F[j(\omega - \omega_0)] + \frac{1}{2} F[j(\omega + \omega_0)] \end{aligned}$$



Example:



$$W_d(j\omega) = \mathcal{F}[w_d(t)] = \int_{-d/2}^{d/2} e^{-j\omega t} dt = \frac{2}{\omega} \sin\left(\frac{\omega d}{2}\right) = \frac{1}{\pi f} \sin \pi f d = d \operatorname{sinc} \pi f d$$

$$F(j\omega) = \mathcal{F}[w_d(t) \cos \omega_0 t] = \frac{\sin \frac{d}{2}(\omega - \omega_0)}{\omega - \omega_0} + \frac{\sin \frac{d}{2}(\omega + \omega_0)}{\omega + \omega_0}$$


Example



$$W_d(j\omega) = F[w_d(t)] = \int_{-d/2}^{d/2} e^{-j\omega t} dt = \frac{2}{\omega} \sin\left(\frac{\omega d}{2}\right)$$

$$F(j\omega) = F[w_d(t) \cos \omega_0 t] = \frac{\sin \frac{d}{2} (\omega - \omega_0)}{\omega - \omega_0} + \frac{\sin \frac{d}{2} (\omega + \omega_0)}{\omega + \omega_0}$$

Example:



$$f(t) = \frac{\sin at}{\pi t} \quad F(j\omega) = ?$$

Sol)

$$W_d(j\omega) = \frac{2}{\omega} \sin\left(\frac{\omega d}{2}\right)$$

$$\rightarrow F[W_d(jt)] = F\left[\frac{2}{t} \sin\left(\frac{td}{2}\right)\right] = 2\pi w_d(-\omega)$$

$$\rightarrow F[f(t)] = F\left[\frac{\sin at}{\pi t}\right] = w_{2a}(-\omega) = \begin{cases} 0 & \omega < |a| \\ 1 & \omega > |a| \end{cases}$$

Fourier Transform of $f'(t)$

$$f(t) \xleftrightarrow{\mathcal{F}} F(j\omega) \text{ and } \lim_{t \rightarrow \pm\infty} f(t) = 0$$



$$f'(t) \xleftrightarrow{\mathcal{F}} j\omega F(j\omega)$$

Pf)

$$\begin{aligned} \mathcal{F}[f'(t)] &= \int_{-\infty}^{\infty} f'(t) e^{-j\omega t} dt \\ &= f(t) e^{-j\omega t} \Big|_{-\infty}^{\infty} + j\omega \int_{-\infty}^{\infty} f(t) e^{-j\omega t} dt \\ &= j\omega F(j\omega) \end{aligned}$$

Fourier Transform of $f^{(n)}(t)$

$$f(t) \xleftrightarrow{\mathcal{F}} F(j\omega) \text{ and } \lim_{t \rightarrow \pm\infty} f(t) = 0$$



$$f^{(n)}(t) \xleftrightarrow{\mathcal{F}} (j\omega)^n F(j\omega)$$

Proved by yourselves

Fourier Transform of $f^{(n)}(t)$

$$f(t) \xleftrightarrow{\mathcal{F}} F(j\omega) \text{ and } \lim_{t \rightarrow \pm\infty} f(t) = 0$$



$$f^{(n)}(t) \xleftrightarrow{\mathcal{F}} (j\omega)^n F(j\omega)$$

Proved by yourselves

Fourier Transform of Integral

$$f(t) \xleftrightarrow{\mathcal{F}} F(j\omega) \text{ and } \int_{-\infty}^{\infty} f(t)dt = F(0) = 0$$


$$\mathcal{F}\left[\int_{-\infty}^t f(x)dx\right] = \frac{1}{j\omega} F(j\omega)$$

Let $\phi(t) = \int_{-\infty}^t f(x)dx$  $\lim_{t \rightarrow \infty} \phi(t) = 0$

$$\mathcal{F}[\phi'(t)] = \mathcal{F}[f(t)] = F(j\omega) = j\omega\Phi(j\omega)$$

$$\Phi(j\omega) = \frac{1}{j\omega} F(j\omega)$$

The Derivative of Fourier Transform

$$\mathcal{F}[-jtf(t)] \longleftrightarrow \frac{dF(j\omega)}{d\omega}$$

Pf)

$$F(j\omega) = \int_{-\infty}^{\infty} f(t)e^{-j\omega t} dt$$


$$\frac{dF(j\omega)}{d\omega} = \frac{d}{d\omega} \int_{-\infty}^{\infty} f(t)e^{-j\omega t} dt = \int_{-\infty}^{\infty} f(t) \frac{\partial}{\partial \omega} e^{-j\omega t} dt$$

$$= \int_{-\infty}^{\infty} [-jtf(t)]e^{-j\omega t} dt = \mathcal{F}[-jtf(t)]$$

Continuous-Time Fourier Transform

Convolution

Basic Concept




$$f_i(t) = a_1 f_{i1}(t) + a_2 f_{i2}(t) \Rightarrow f_o(t) = L[a_1 f_{i1}(t) + a_2 f_{i2}(t)]$$


A linear system satisfies

$$f_o(t) = a_1 L[f_{i1}(t)] + a_2 L[f_{i2}(t)]$$
$$= a_1 f_{o1}(t) + a_2 f_{o2}(t)$$

Basic Concept



Basic Concept






A causal system satisfies


$$f_i(t) = 0 \text{ for } t < t_0 \quad \longrightarrow \quad f_o(t) = 0 \text{ for } t < t_0$$

Which of the following systems are causal?

Basic Concept



Unit Impulse Response




Facts: $\int_{-\infty}^{\infty} f(\tau)\delta(t-\tau)d\tau = \int_{-\infty}^{\infty} f(t-\tau)\delta(\tau)d\tau = f(t)$

$\rightarrow L[f(t)] = L\left[\int_{-\infty}^{\infty} f(\tau)\delta(t-\tau)d\tau\right] = \int_{-\infty}^{\infty} f(\tau)L[\delta(t-\tau)]d\tau$

$= \int_{-\infty}^{\infty} f(\tau)h(t-\tau)d\tau$ **Convolution**


Unit Impulse Response



$$L[f(t)] = f(t) * h(t)$$

$$= \int_{-\infty}^{\infty} f(\tau) h(t - \tau) d\tau \quad \text{Convolution}$$

Unit Impulse Response



Convolution Definition

The **convolution** of two functions $f_1(t)$ and $f_2(t)$ is defined as:

$$\begin{aligned} f(t) &= \int_{-\infty}^{\infty} f_1(\tau) f_2(t - \tau) d\tau \\ &= f_1(t) * f_2(t) \end{aligned}$$


Properties of Convolution

$$f_1(t) * f_2(t) = f_2(t) * f_1(t)$$

$$\begin{aligned} f_1(t) * f_2(t) &= \int_{-\infty}^{\infty} f_1(\tau) f_2(t - \tau) d\tau = \int_{\tau=-\infty}^{\tau=\infty} f_1(\tau) f_2(t - \tau) d\tau \\ &= \int_{t-\tau=-\infty}^{t-\tau=\infty} f_1(t - \tau) f_2[t - (t - \tau)] d(t - \tau) \\ &= - \int_{\tau=\infty}^{\tau=-\infty} f_1(t - \tau) f_2(\tau) d\tau \\ &= \int_{-\infty}^{\infty} f_1(t - \tau) f_2(\tau) d\tau = f_2(t) * f_1(t) \end{aligned}$$

Properties of Convolution

$$f_1(t) * f_2(t) = f_2(t) * f_1(t)$$



Properties of Convolution



$$[f_1(t) * f_2(t)] * f_3(t) = f_1(t) * [f_2(t) * f_3(t)]$$

Prove by yourselves


The following two systems are identical

Properties of Convolution

$$[f_1(t) * f_2(t)] * f_3(t) = f_1(t) * [f_2(t) * f_3(t)]$$




Properties of Convolution



$$\begin{aligned} f(t) * \delta(t) &= \int_{-\infty}^{\infty} f(\tau) \delta(t - \tau) d\tau \\ &= \int_{-\infty}^{\infty} f(t - \tau) \delta(\tau) d\tau \\ &= f(t) \end{aligned}$$

Properties of Convolution

$$f(t) * \delta(t) = f(t)$$





A block diagram illustrating the identity property of convolution. On the left, the equation $f(t) * \delta(t) = f(t)$ is shown in a cyan box. To the right, a blue $f(t)$ is connected by a red arrow to a yellow rectangular block labeled $\delta(t)$. Another red arrow connects the block to a final blue $f(t)$.

$$f(t) * \delta(t - T) = f(t - T)$$

$$\begin{aligned} f(t) * \delta(t - T) &= \int_{-\infty}^{\infty} f(\tau) \delta(t - T - \tau) d\tau \\ &= \int_{-\infty}^{\infty} f(t - T - \tau) \delta(\tau) d\tau \\ &= f(t - T) \end{aligned}$$




Properties of Convolution

$$f(t) * \delta(t - T) = f(t - T)$$



System function $\delta(t-T)$ serves as an *ideal delay* or a *copier*.

$$f(t) * \delta(t-T) = f(t-T)$$



Properties of Convolution

$$f_1(t) * f_2(t) \xleftrightarrow{F} F_1(j\omega)F_2(j\omega)$$

$$\begin{aligned} F[f_1(t) * f_2(t)] &= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} f_1(\tau) f_2(t - \tau) d\tau \right] e^{-j\omega t} dt \\ &= \int_{-\infty}^{\infty} f_1(\tau) \left[\int_{-\infty}^{\infty} f_2(t - \tau) e^{-j\omega t} dt \right] d\tau \\ &= \int_{-\infty}^{\infty} f_1(\tau) F_2(j\omega) e^{-j\omega \tau} d\tau \\ &= F_2(j\omega) \int_{-\infty}^{\infty} f_1(\tau) e^{-j\omega \tau} d\tau = F_1(j\omega) F_2(j\omega) \end{aligned}$$

Time Domain

convolution

Frequency Domain

multiplication

$$f_1(t) * f_2(t) \xleftrightarrow{F} F_1(j\omega)F_2(j\omega)$$

$$\begin{aligned} F[f_1(t) * f_2(t)] &= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} f_1(\tau) f_2(t - \tau) d\tau \right] e^{-j\omega t} dt \\ &= \int_{-\infty}^{\infty} f_1(\tau) \left[\int_{-\infty}^{\infty} f_2(t - \tau) e^{-j\omega t} dt \right] d\tau \\ &= \int_{-\infty}^{\infty} f_1(\tau) F_2(j\omega) e^{-j\omega \tau} d\tau \\ &= F_2(j\omega) \int_{-\infty}^{\infty} f_1(\tau) e^{-j\omega \tau} d\tau = F_1(j\omega) F_2(j\omega) \end{aligned}$$



Time Domain

Frequency Domain

convolution

multiplication

$$f_1(t) * f_2(t) \xleftrightarrow{F} F_1(j\omega)F_2(j\omega)$$




Time Domain

convolution

Frequency Domain


multiplication

$$f_1(t) * f_2(t) \xleftrightarrow{F} F_1(j\omega)F_2(j\omega)$$



Properties of Convolution


$$f_1(t) * f_2(t) \xleftrightarrow{F} F_1(j\omega)F_2(j\omega)$$



An Ideal Low-Pass Filter

Properties of Convolution

$$f_1(t) * f_2(t) \xleftrightarrow{F} F_1(j\omega)F_2(j\omega)$$



An Ideal High-Pass Filter

Properties of Convolution

$$f_1(t) f_2(t) \xleftrightarrow{\mathcal{F}} \frac{1}{2\pi} \int_{-\infty}^{\infty} F_1(j\theta) F_2[j(\omega - \theta)] d\theta$$

$$f_1(t) f_2(t) \xleftrightarrow{\mathcal{F}} \frac{1}{2\pi} F_1(j\omega) * F_2(j\omega)$$

Prove by yourselves

Time Domain

multiplication

Frequency Domain

convolution

$$f_1(t)f_2(t) \xleftrightarrow{\mathcal{F}} \frac{1}{2\pi} \int_{-\infty}^{\infty} F_1(j\theta)F_2[j(\omega - \theta)]d\theta$$

$$f_1(t)f_2(t) \xleftrightarrow{\mathcal{F}} \frac{1}{2\pi} F_1(j\omega) * F_2(j\omega)$$

Prove by yourselves


Continuous-Time Fourier Transform


Parseval's Theorem

Properties of Convolution

$$\int_{-\infty}^{\infty} [f_1(t) f_2(t)] dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} F_1(j\omega) F_2[-j\omega] d\omega$$

$$f_1(t) f_2(t) \xleftrightarrow{\mathcal{F}} \frac{1}{2\pi} \int_{-\infty}^{\infty} F_1(j\theta) F_2[j(\omega - \theta)] d\theta$$



$$\int_{-\infty}^{\infty} [f_1(t) f_2(t)] e^{j\omega t} dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} F_1(j\theta) F_2[j(\omega - \theta)] d\theta$$


$$\int_{-\infty}^{\infty} [f_1(t) f_2(t)] dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} F_1(j\omega) F_2[j(-\omega)] d\omega$$

Properties of Convolution


$$\int_{-\infty}^{\infty} [f_1(t) f_2(t)] dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} F_1(j\omega) F_2[-j\omega] d\omega$$

If $f_1(t)$ and $f_2(t)$ are real functions,



$$\int_{-\infty}^{\infty} [f_1(t) f_2(t)] dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} F_1(j\omega) F_2^*[j\omega] d\omega$$

$f_2(t)$ real



$$F_2[-j\omega] = F_2^*[j\omega]$$

Parseval's Theorem: Energy Preserving

$$\int_{-\infty}^{\infty} |f(t)|^2 dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} |F(j\omega)|^2 d\omega$$

$$F[f^*(t)] = \int_{-\infty}^{\infty} f^*(t) e^{-j\omega t} dt = \left(\int_{-\infty}^{\infty} f(t) e^{j\omega t} dt \right)^* = F^*(-j\omega)$$

$$\int_{-\infty}^{\infty} |f(t)|^2 dt = \int_{-\infty}^{\infty} f(t) f^*(t) dt$$

$$= \frac{1}{2\pi} \int_{-\infty}^{\infty} F(j\omega) F^*[-(-j\omega)] d\omega = \frac{1}{2\pi} \int_{-\infty}^{\infty} |F(j\omega)|^2 d\omega$$